

Ly-49W, an Activating Receptor of Nonobese Diabetic Mice With Close Homology to the Inhibitory Receptor Ly-49G, Recognizes H-2D^k and H-2D^d 1

Elizabeth T. Silver, Dong-Er Gong, Bart Hazes, and Kevin P. Kane²

The diversity and ligand specificity of activating Ly-49 receptors expressed by murine NK cells are largely unknown. We cloned a new Ly-49-activating receptor, expressed by NK cells of the nonobese diabetic mouse strain, which we have designated Ly-49W. Ly-49W is highly related to the known inhibitory receptor Ly-49G in its carbohydrate recognition domain, exhibiting 97.6% amino acid identity in this region. We demonstrate that the 4D11 and Cwy-3 Abs, thought to be Ly-49G specific, also recognize Ly-49W. Rat RNK-16 cells transfected with Ly-49W mediated reverse Ab-dependent cellular cytotoxicity of FcR-positive target cells, indicating that Ly-49W can activate NK-mediated lysis. We further show that Ly-49W is allo-MHC specific: Ly-49W transfectants of RNK-16 only lysed Con A blasts expressing H-2^k or H-2^d haplotypes, and Ab-blocking experiments indicated that H-2D^k and D^d are ligands for Ly-49W. Ly-49W is the first activating Ly-49 receptor demonstrated to recognize an H-2^k class I product. Ly-49G and Ly-49W represent a new pair of NK receptors with very similar ligand-binding domains, but opposite signaling functions. *The Journal of Immunology*, 2001, 166: 2333–2341.

Natural killer cells function as a first line of defense to eliminate tumor cells or virally infected cells without prior sensitization (1, 2). NK cells also mediate allogeneic bone marrow rejection as well as hybrid resistance, wherein an F₁ mouse rejects a bone marrow transplant from a parent (3, 4). The cytotoxicity of NK cells appears to be regulated by opposing signals generated upon interaction with potential target cells. Positive signals are induced upon the interaction of activating receptor(s) with largely undefined target cell surface ligands (5–7), while NK activation is negatively regulated when inhibitory receptors bind class I MHC molecules on the target cell (8). NK cells can lyse cells with deficient self class I expression, but spare cells that express normal levels of self class I MHC molecules. Such observations have resulted in the “missing self” hypothesis (9), which holds as one of its tenets an *in vivo* selection process in which each mature NK cell will express at least one inhibitory receptor recognizing self class I MHC to ensure NK cell self-tolerance (9, 10).

NK cells express inhibitory receptors from the Ig- and the C-type lectin superfamilies, with the human killer Ig-related receptor (KIR)³ and the murine Ly-49 families being the most highly characterized of the Ig- and lectin-like NK-inhibitory receptors, respec-

tively (11). While KIR and Ly-49 are structurally dissimilar, they are functionally equivalent, serving as inhibitory receptors that can distinguish class I alleles in their MHC recognition. A common feature of inhibitory receptors is the presence of an immunoreceptor tyrosine-based inhibitory motif (ITIM) (12). Engagement of the inhibitory receptor with its class I ligand on the target cell results in tyrosine phosphorylation of the ITIM (13, 14). The phosphorylated ITIM recruits the tyrosine phosphatase SHP-1, which then dephosphorylates plasma membrane-proximal targets in the NK activation cascade, thus blocking NK activation (13, 14).

All NK cell-inhibitory receptor families contain members that lack ITIM sequences. The ITIM-lacking molecules have a charged residue in their transmembrane domains to allow interaction with signaling adaptor molecules (12). For the KIR and Ly-49 families, this adaptor is the DAP12 signaling molecule that contains an immunoreceptor tyrosine-based activation motif (11, 15–17). DAP12 interacts with the ITIM-lacking receptors via a noncovalent interaction in the transmembrane domain (11, 17). Ligation of the receptor recruits and activates Syk kinase, leading to subsequent downstream activation events, target cell cytolysis, and cytokine production (18–20). Some Ly-49 proteins can be grouped into inhibitory/activating pairs based on high sequence identities in their external domains (12). The functional significance of these pairs is unknown, but the considerable resemblance suggests that they recognize the same ligands. Ly-49D and Ly-49P are the only activating Ly-49 receptors for which a ligand has been determined. Ly-49D and Ly-49P recognize the H-2D^d class I MHC molecule, similar to the ITIM-containing inhibitory receptors, Ly-49A and Ly-49G (21–25).

Studies of Ly-49 receptor function have largely been confined to the C57BL/6 (B6) mouse strain. In this study, we examined Ly-49 gene expression and function in the nonobese diabetic (NOD) mouse, to explore strain-to-strain variation in Ly-49 expression, and because of its well-established immune dysregulation and spontaneous onset of autoimmune diabetes (26). We describe Ly-49W, a novel activating Ly-49 receptor expressed in NOD mice with high homology in its external domain to the inhibitory Ly-49G receptor. We also demonstrate that Ly-49W recognizes class

Department of Medical Microbiology and Immunology, University of Alberta, Edmonton, Alberta, Canada

Received for publication July 10, 2000. Accepted for publication November 28, 2000.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ This work was supported by an operating grant from the National Cancer Institute of Canada and the Canadian Cancer Society. E.T.S. is supported by an Alberta Heritage Foundation for Medical Research studentship. K.P.K. is an Alberta Heritage Foundation for Medical Research senior scholar.

² Address correspondence and reprint requests to Dr. Kevin P. Kane, Department of Medical Microbiology and Immunology, 660 HMRC, University of Alberta, Edmonton, Alberta, Canada T6G 2S2. E-mail address: kevin.kane@ualberta.ca

³ Abbreviations used in this paper: KIR, killer Ig-related receptor; ADCC, Ab-dependent cellular cytotoxicity; CRD, carbohydrate recognition domain; ITIM, immunoreceptor tyrosine-based inhibitory motif; NOD, nonobese diabetic; NOR, nonobese diabetes-resistant; PA, protein A; PG, protein G; rADCC, reverse ADCC.

I MHC molecules with strong and moderate reactivity toward H-2D^k and H-2D^d, respectively. This is the first report of an activating Ly-49 molecule recognizing an H-2^k product and provides further support for the possibility that activating Ly-49 receptors recognize class I MHC ligands.

Materials and Methods

Animals

Five- to 8-wk-old female AKR/J (H-2^k), CBA/J (H-2^k), BALB/c (H-2^d), DBA/2J (H-2^d), C57BL/6 (H-2^b), NOD (H-2K^d,D^b), nonobese diabetes-resistant (NOR) (H-2K^d,D^b), B10.BR (H-2^b), B10.D2 (B10.D2-H2^d/nSn; H-2^d), B10.S (H-2^s), B10 (C57BL/10; H-2^b), B10.d/b (B10.HTG-H2^s/2Cy; H-2K^d,D^b), and B10.b/d (B10.A-H2^b/H2-T18^a(5R)/SgSn; H-2K^b,D^d) mice were purchased from The Jackson Laboratory (Bar Harbor, ME). Additional 5- to 8-wk-old NOR and NOD mice were obtained from the University of Alberta (Edmonton, Alberta, Canada) mouse-breeding facility.

Hybridomas and Abs

Hybridomas producing the following Abs: 4D11 (rat IgG2a), anti-Ly-49G (27); Cwy-3 (IgG1), anti-Ly-49G (28); M1/42 (rat IgG2a), anti-mouse class I MHC (29); 34-5-8S (IgG2a), anti-H-2D^d α 1/ α 2 domain epitope (30); 34-2-12S (IgG2a) anti-H-2D^d α 3 domain epitope (30); B8-24-3 (IgG1), anti-H-2K^b (31); B27 M1 (IgG2a), anti-HLA-B27, B7 (32); and BB7.1 (IgG1), anti-HLA-B7 (33) were obtained from American Type Culture Collection (Manassas, VA), except Cwy-3, which was generated in this laboratory. Abs were prepared by ammonium sulfate precipitation, and PBS dialysis of tissue culture supernatants obtained from hybridomas grown in protein-free hybridoma medium. Purified OX-8 (IgG1) anti-rat CD8 α (34), 11-4.1 (IgG2a) anti-K^k (35), and 15-5-5S (IgG2a) anti-D^k (36) Abs were purchased from BD Pharmingen (San Diego, CA). Purified rat IgG was purchased from Sigma-Aldrich (Oakville, Canada). FITC-coupled rat anti-mouse IgG, goat anti-rat IgG, and mouse anti-rat IgG were purchased from Jackson ImmunoResearch (West Grove, PA).

Cell lines

RNK-16, a spontaneous F344 rat strain NK cell leukemia cell line (37), was kindly provided by Dr. Mary Nakamura at the University of California at San Francisco. The RNK-16 cells were maintained in RPMI 1640 supplemented with 10% FCS, L-glutamine, penicillin, streptomycin, and 5×10^{-5} M 2-ME (RNK medium). COS-7 SV40-transformed African green monkey kidney cells were provided by Dr. John Elliott (University of Alberta). COS-7 cells were grown in Opti-MEM I medium (Life Technologies, Burlington, Ontario, Canada), containing 4% heat-inactivated FCS (Mediatech, Montreal, Canada) and 5×10^{-5} M 2-ME.

NK cell propagation, RNA preparation, and cDNA synthesis

IL-2-activated NK cells were prepared from NOD, NOR, or C57BL/6 strain spleen cells harvested from 6- to 8-wk-old female mice, as described by Smith and coworkers (38). Total cellular RNA was isolated with TRIzol reagent (Life Technologies) and reverse transcribed using SuperScript II (Life Technologies) with an oligo(dT) primer.

Cloning of NOD Ly-49 transcripts

Ly-49 transcripts were amplified with Advantage cDNA Polymerase (Clontech, Palo Alto, CA) using the sense cloning primer 5'-CCCAAGAT GAGTGAGCAGGAGG and the antisense cloning primer 5'-GAGAGT CAATGAGGGAATTTATCC. The PCR products were purified using QIAquick spin columns (Qiagen, Santa Clarita, CA) and directly ligated into the TA cloning vector p3T (Miltenyi Biotec, Göttingen, Germany). The individual cDNA clones were divided into two groups: ITIM-encoding clones, which could be PCR amplified using the antisense cloning primer, and the sense primer 5'-GGAGACTCAAGGGCCAGAG, or ITIM-lacking clones, which could be PCR amplified by using the antisense cloning primer and the sense primer 5'-CTRRAAAAGCTGGCCTCAGAGTY (in which R = A/G and Y = C/T). To verify the nucleotide sequences of the ends of the open reading frames and to obtain sequence of the noncoding gene-flanking regions for cloning, we performed 5' rapid amplification of cDNA ends of Ly-49W. Poly(A)⁺ RNA was isolated from total RNA, obtained from IL-2-activated NK cells using Oligotex spin columns (Qiagen), and used to synthesize a cDNA library with the Marathon cDNA Amplification Kit (Clontech). The gene-specific primer used for Ly-49W 5' rapid amplification of cDNA ends was 5'-GCCTGGCCTACTCTATA GATGTAGAAC. To obtain full-length cDNA clones, we used the follow-

ing primers: Ly-49G, 5'-ATTTAACTGAGAACA TACTTCATACAT CAT (sense) and 5'-GATTGTCTCTCTTTGCACTTTTACA (antisense); Ly-49W, 5'-TTTAAAAGAGAACATACTACTACATCCTC (sense) and the same antisense primer as Ly-49G. Sequencing reactions were performed using a dideoxy terminator method and analyzed on an ABI 373A automated sequencer (Applied Biosystems, Foster City, CA).

Transfection and flow cytometric analysis

COS-7 cells. DNA containing the coding regions of Ly-49 cDNAs was inserted into the *Xba*I/*Eco*RI sites of the mammalian expression vector pCI-neo (Promega, Madison, WI). Sequence encoding the mature mouse DAP12 protein was inserted into the pFLAG-CMV-1 expression vector (Sigma-Aldrich) to create an epitope-tagged chimeric protein. Vectors were then transfected into COS cells using LipofectAMINE (Life Technologies). Approximately 48 h after transfection, the cells were incubated with the mAbs 4D11 or Cwy-3, or isotype control Abs M1/42 or B8-24-3, respectively. Secondary Abs, either FITC-labeled goat anti-rat IgG or rat anti-mouse IgG, respectively, were subsequently added for an additional incubation, whereupon the samples were analyzed on a FACScan flow cytometer (Becton Dickinson, Mountain View, CA).

RNK cells. The cDNA encoding Ly-49W was inserted into the *Xho*I/*Xba*I sites of the bicistronic vector BSR α EN (generously provided by Dr. Andrew Shaw, Washington University, St. Louis, MO) and transfected into RNK-16 cells using the protocol described by Nakamura et al. (14). In brief, four million cells were transfected with 20 μ g of plasmid linearized with *Sac*I by electroporation at 200 mV and 960 μ FD. Transfected cells were cloned in 96-well microtiter plates in complete RNK medium supplemented with 1 mg/ml G418 for drug selection. Expression of Ly-49W on transfectant clones was measured by FACS analysis with the 4D11 Ab after blocking FcR with normal mouse serum. Reactivity of transfectant clones with Cwy-3 and OX-8 Abs was also determined after blocking with purified rat IgG. The BB7.1 Ab was used as an isotype control. Primary Ab binding was detected with mouse anti-rat and rat anti-mouse fluorescein-coupled Abs, respectively, using a FACScan flow cytometer.

Generation of Con A T cell blast target cells

Con A-activated T cell blasts were prepared from spleen cells of various mouse strains. Fifteen million spleen cells were cultured at 5×10^6 cells/ml in RPMI 1640 with 10% heat-inactivated FCS, 2-ME, and 3 μ g/ml Con A (Sigma-Aldrich) for 48 h. Blast cells were recovered after three washes in RPMI 1640 medium.

Cytotoxicity assays

Target cells were labeled at 37°C with 100–150 μ Ci of Na⁵¹CrO₄ (⁵¹Cr) (Mandel/NEN Life Science Products, Guelph, Canada), for 1 h if tumor cells, or 1.5 h if Con A blast cells. Following extensive washes, 1×10^4 ⁵¹Cr-labeled target cells were incubated for 4–5 h at 37°C in V-bottom microtiter plates with RNK-16 cells or RNK-16 cells transfected with Ly-49W at various E:T ratios in triplicate. After the incubation, plates were centrifuged for 5 min, and 100 μ l of supernatant was removed and counted in a gamma counter. The percent specific lysis was determined as (experimental release – spontaneous release)/(maximum release – spontaneous release) \times 100. To perform the reverse Ab-dependent cellular cytotoxicity (rADCC) experiments, untransfected RNK-16 cells and the Ly-49W transfectants of RNK-16 were preincubated for 15 min with 20 μ g/ml of the Cwy-3 or OX-8 Abs or medium alone before addition of FcR-expressing YB2/0 target cells and subsequent 4-h cytotoxic assay, as described. For Ab inhibition experiments, Abs were incubated with soluble protein A (PA) (2 μ g/10 μ g of mAb; Sigma-Aldrich) or a mix of PA and protein G (PG) (4 μ g/10 μ g of mAb; ICN Pharmaceuticals, Costa Mesa, CA) for 30 min before addition to effector cells or target cells. Effector cells or target cells were preincubated with the mAb and PA for 15 min before the cytotoxicity assay. In the case of two Abs being employed simultaneously in the same wells in receptor-blocking studies, each Ab is used at the indicated concentrations in the figure. The mAbs and PA or PA/PG were present throughout the cytotoxicity assays. All cytotoxicity assays were repeated a minimum of three times.

Results

Cloning of noninhibitory Ly-49 members with homology to Ly-49G

Recent studies suggest that Ly-49 gene expression varies between mouse strains (25, 39, 40). Furthermore, the complex hybridization patterns in Southern blots of various mouse strains suggest that not

all Ly-49 family members have been identified (41). Indeed, investigation of Ly-49 expression in mouse strains such as 129J and CBA/J have resulted in the identification of novel activating Ly-49 molecules (40, 42). In this study, we examine Ly-49 expression in the NOD mouse strain, an animal model of insulin-dependent diabetes mellitus. NOR strain mice are identical with NOD at most, but not all, genetic loci and do not develop diabetes. We designed an RT-PCR strategy to clone cDNAs encoding both activating and inhibitory Ly-49 family members from IL-2-activated NK cells of NOD and NOR strain mice. From this, we obtained cDNAs encoding both ITIM-containing and ITIM-lacking receptors.

The ITIM-containing cDNAs prepared from NOD IL-2-activated NK cells include a novel allele of the inhibitory *Ly-49G* gene with a nucleotide identity of 98.8% and 98.9%, and an amino acid identity of 97% and 98.1% compared with the C57BL/6 and BALB/c alleles, respectively (Fig. 1A). The NOD *Ly-49G* allele more closely resembles both the C57BL/6 and BALB/c alleles than they resemble each other (97.9% nucleotide and 95.9% amino acid identities). The NOD transcripts described in this comparison correspond to *Ly-49G2*, first defined in the C57BL/6 (B6) strain (Fig. 1A). NOR NK cells also express *Ly-49G2* (GenBank accession number AF074457) with 100% sequence identity to NOD *Ly-49G2*. Similar to the B6 strain, NOD NK cells also express alternatively spliced RNA transcripts encoding a larger *Ly-49G* form, *Ly-49G1^{NOD}*, in which there is an extension of 13 aa residues in the extracellular membrane-proximal stalk domain of the receptor (GenBank accession number AF283248). In addition, NOD NK cells express *Ly-49G* transcripts that, through alternative splicing, lack exon 3 encoding the transmembrane segment and a portion of the cytoplasmic domain (GenBank accession number AF283253).

The ITIM-lacking NOD IL-2-activated NK cDNAs include sequences of two novel *Ly-49* transcripts. Both sequences show strong similarity with two genomic fragments from B6 mice that were originally designated *Ly-49M* (43). To determine the relationship of our new sequences with *Ly-49M* from the B6 strain, we cloned and sequenced the full *Ly-49M* transcript from B6 mouse cDNA (Fig. 1B). This showed that *Ly-49M* is not functionally expressed in B6 mice as there is a premature stop codon near the predicted beginning of exon 4 (Fig. 1B). The B6 *Ly-49M* sequence differs by only 10 bases and 8 aa from one of our new sequences, corresponding to 98.7% and 97% sequence identity on the DNA and protein levels, respectively. This extent of sequence identity is comparable with that found between alleles of *Ly-49G*. Thus, we are tentatively designating this NOD *Ly-49* product as the NOD allele of *Ly-49 m*. Interestingly, the NOD gene does not contain the premature stop codon, suggesting that it represents a functional receptor. This is an additional example of variation in *Ly-49* expression between different mouse strains.

Our other new NOD *Ly-49* transcript also resembles *Ly-49M* with 18 bases and 15 aa substituted, corresponding to 97.7% and 94.3% sequence identity on the DNA and protein level, respectively. However, this product must represent a distinct gene because we have already shown above that NOD encodes an even more closely related *Ly-49M* homologue. Hence, we have designated this NOD *Ly-49* transcript *Ly-49W* (Fig. 1A). There are two distinct *Ly-49W* mRNA transcripts that result from alternative splicing at the beginning of exon 3. We have previously demonstrated that this form of alternative splicing also occurs in *Ly-49D* and *Ly-49H* (44). Splice variants containing the coding sequence for cytoplasmic residues Val-Cys-Ser are named *Ly-49D1* and *Ly-49H1*, while those lacking this coding sequence are termed *Ly-49D2* and *Ly-49H2*. Hence, we are using the designations *Ly-49W1* and *Ly-49W2* for transcripts that encode or lack the Val-Cys-Ser sequence, respectively (Fig. 1B). We have also found




FIGURE 1. Comparison of the predicted amino acid sequences of Ly-49W with Ly-49G, Ly-49L, and Ly-49M. Dashes indicate sequence identity, asterisks indicate sequence gaps, and potential N-linked glycosylation sites (shown in A) are boxed (unshaded). Predicted protein domains are indicated by horizontal arrows above the sequences. A, The predicted amino acid sequence of Ly-49W is aligned with those of the Ly-49G alleles from the C57BL/6, BALB/c, and NOD mouse strains. The ITIM sequence is shaded and boxed with dotted lines. B, The alignment of the predicted protein sequences of splice variants Ly-49W1 and Ly-49W2 with Ly-49L2 of BALB/c mice as well as Ly-49M from C57BL/6 and NOD mouse strains. The arginine in the transmembrane region is shaded and boxed. The termination codon is indicated by a lower case bold z. Nucleotide sequences are available under the GenBank accession numbers: AF283250 (*Ly-49W1^{NOD}*), AF283251 (*Ly-49W2^{NOD}*), AF307946 (*Ly-49G2^{BALB/c}*), AF283249 (*Ly-49G2^{NOD}*), AF307947 (*Ly-49L^{BALB/c}*), AF028133 (*Ly-49M^{B6}* (partial, exons 4 and 7)), AF074460 (*Ly-49M^{B6}*), and AF283252 (*Ly-49M^{NOD}*).

both splice variants with identical sequences in the NOR mouse strain (GenBank accession numbers AF074459 and AF074463).

Recently, the complete sequence of a novel receptor, Ly-49L, was determined. Ly-49L is found in CBA/J, C3H (42), and BALB/c mice (Fig. 1B), and its nucleotide and amino acid sequence does not differ in these strains. Ly-49L has 94.3% amino acid sequence identity with Ly-49W (Fig. 1B), the same level of

divergence as observed between Ly-49W and Ly-49M. However, Ly-49W and Ly-49L amino acid sequences are identical in exons 2 to 5, with all substitutions taking place in the C-terminal region encoded by exons 6 and 7 (Fig. 1B, and data not shown). The genetic relationship of Ly-49W and Ly-49L is unclear. The overall level of amino acid identity strongly suggests that Ly-49L and Ly-49W are separate genes. It remains a possibility, however, that Ly-49W and Ly-49L are alleles of the same gene, but a substitution of exons 6 and 7 from another gene has occurred in Ly-49L or Ly-49W. Detailed genomic analysis of Ly-49 genes in multiple mouse strains will be necessary to address these possibilities. In any event, the amino acid sequence of the external domain of Ly-49W is closer to that of Ly-49G than those of Ly-49M or Ly-49L.

The carbohydrate recognition domain (CRD) sequence of Ly-49W (exons 5 to 7) is strikingly similar to the same region of the inhibitory receptor Ly-49G, with an amino acid sequence identity of 97.6% compared with the Ly-49G alleles of both the B6 and NOD strains (Fig. 1A). In the case of the stalk region, these numbers are 83.8% and 82.4%, respectively. Interestingly, the cytoplasmic/transmembrane domain is rather different as it lacks the ITIM motif and contains an arginine in the middle of its transmembrane domain (Fig. 1). This is the hallmark of the known activating receptors Ly-49D, Ly-49H, and Ly-49P (17, 25, 40). Below, we will present evidence that Ly-49W is indeed an activating receptor. Accordingly, Ly-49G/Ly-49W form an inhibitory/activating pair with opposite regulatory effects, but closely related CRDs. This is analogous to other Ly-49 inhibitory/activating pairs that have previously been reported (25, 40, 41). The studies described below will examine the function and specificity of Ly-49W.

Ly-49G-reactive Abs 4D11 and Cwy-3 recognize NOD Ly-49W

Serological determinants may be shared between NOD Ly-49G and Ly-49W due to the high degree of amino acid identity in their external domains. To address this possibility, we transiently expressed cDNAs encoding these molecules in COS cells and performed FACS analysis with Abs reactive with Ly-49G2. The 4D11 Ab was initially demonstrated to detect an epitope expressed on several mouse strains (27) and subsequently was determined to recognize Ly-49G (23). The Cwy-3 mAb has been shown to have a much more restricted specificity by recognizing Ly-49G of C57BL/6, but not several other strains (28). Cwy-3 and 4D11 recognize Ly-49G of NOD origin, as indicated by substantial staining of COS cell transfectants with these Abs (Fig. 2, upper panels). Thus, Cwy-3 recognizes the NOD and B6 alleles of Ly-49G. Ly-49W expression on COS cells is not readily detected with Cwy-3 or 4D11 Abs, unless Ly-49W is cotransfected with DAP12 (Fig. 2, lower panels). Thus, Ly-49W is recognized by the Ly-49G-specific mAbs 4D11 and Cwy-3, but, similar to other noninhibitory Ly-49 family members (17, 40), appears to require the accessory molecule DAP12 for efficient cell surface expression.

Ly-49W mediates rADCC

The noninhibitory Ly-49D, Ly-49H, and Ly-49P receptors have been shown to be activating receptors by their ability to mediate rADCC, or redirected lysis, against FcR-expressing targets (25, 45, 46). Because the Abs that bind Ly-49W also recognize the inhibitory receptor Ly-49G, the experimental system we chose (14) employed the RNK-16 NK cell line transfected with mouse Ly-49W as effector cells. In this way, we could be certain there was no Ab cross-reactivity and the effector cells would be Ly-49W⁺, yet Ly-49G⁻. Transfected RNK-16 cells were screened by FACS analysis to select for clones stably expressing Ly-49W. A number of trans-




FIGURE 2. Ly-49W is recognized by the 4D11 and Cwy-3 Abs. COS-7 cells transfected with Ly-49G from C57BL/6 (B6) or NOD mice, or Ly-49W alone or Ly-49W cotransfected with murine DAP12 were analyzed by flow cytometry using the Abs 4D11 or Cwy-3 (shaded), or the isotype control Abs M1/42 or B8-24-3 (unshaded), respectively.

fectant clones expressing Ly-49W were obtained as determined by acquisition of reactivity with the 4D11 and Cwy-3 Abs, and three clones were selected for study (Fig. 3). Cotransfection of mouse DAP12 was not required for expression of Ly-49W on RNK-16 cells as is observed with Ly-49D, another noninhibitory Ly-49 (24), presumably because rat DAP-12 substitutes for the mouse accessory protein. Expression of rat CD8 α , a constitutively expressed endogenous rat receptor, was retained following transfection with Ly-49W (Fig. 3).




FIGURE 3. Expression of Ly-49W on RNK-16 transfectants. FACS analysis of Ly-49W-transfected RNK-16 clones 2C4, 7E8, and 10G5 with the Abs 4D11 or Cwy-3 (shaded), or no first Ab, or BB7.1 isotype control (unshaded), respectively. Expression of rat CD8 α on RNK-16 cells was analyzed with the OX-8 Ab (shaded) or the isotype control BB7.1 (unshaded). Ab binding was detected with fluorescein-coupled mouse anti-rat or rat anti-mouse Abs.

RNK-16 cells can efficiently lyse the FcR-expressing target cell line YB2/0 (14) (Fig. 4A), but there is often a marked decrease in target cell killing when the RNK-16 cells are transfected with an activating receptor, such as Ly-49D (24). This effect may be due to a competition for, and sequestration of, a limiting activation component, possibly DAP12, by the transfected activating receptor, relative to an activating receptor that is normally involved in YB2/0 recognition. Such a substantial reduction of direct YB2/0 cytotoxicity is seen with two Ly-49W transfected clones, 2C4 and 10G5 (Fig. 4, B and C). Lysis of FcR-expressing YB2/0 target cells by the Ly-49W-transfected RNK-16 clones 2C4 and 10G5 could be substantially increased when Ly-49W on the effector cells was cross-linked by the Cwy-3 Ab (Fig. 4, B and C). In contrast, cross-linking of effector cell CD8 α by the OX-8 Ab did not increase target cell lysis, although there is comparable cell surface expression of CD8 α and Ly-49W on each of the transfected RNK clones tested (Fig. 3). These results indicate that Ly-49W can mediate rADCC and can be considered an activating receptor. Because high baseline levels of YB2/0 lysis appear to be retained for the Ly-49W transfectant clone 7E8, perhaps due to a greater abundance of signaling molecules, little augmentation of cytotoxicity by Cwy-3 cross-linking of Ly-49W expressed on 7E8 can be detected with this clone (Fig. 4D). Untransfected RNK-16-mediated lysis of YB2/0 cells was unaffected by the Cwy-3 and OX-8 Abs (Fig. 4A).

The cytolytic activity of Ly-49W-transfected RNK-16 cells is MHC dependent

The H-2D^d class I MHC molecule is a ligand for some inhibitory Ly-49 receptors, including Ly-49G (23) and the activating receptor Ly-49D (24). To test whether class I MHC molecules, especially of the H-2^d haplotype, are recognized by Ly-49W, we examined Ly-49W-mediated killing of Con A-activated T cell blast target cells prepared from splenocytes of mouse strains expressing different MHC haplotypes. None of the Ly-49W transfectant clones lysed B6 Con A targets (H-2^b), whereas all of the transfectants lysed Con A blasts from the AKR and CBA mouse strains expressing H-2^k MHC molecules (Fig. 5A). The B10.D2 and DBA/2 mouse strains bearing H-2^d MHC molecules were lysed at a very

low level except in the case of the highly lytic clone 7E8, which kills H-2^d Con A blasts at a moderate level (Fig. 5A). No lysis of any of the Con A T cell blast targets was observed with the untransfected RNK-16 cells. Together, these results suggest that Ly-49W may recognize H-2^k, and possibly H-2^d, product(s).

To further define the importance of MHC molecules in Ly-49W recognition, we examined the lysis of Con A blasts generated from congenic B10 mouse strains that differ only in their MHC genes, by the Ly-49W transfectants. Striking differences were observed in the lysis of the Con A blasts generated from the B10 congenic mice. The B10.BR (H-2^k) Con A blasts were readily lysed by all three Ly-49W transfectant clones (Fig. 5B). The B10.D2 (H-2^d) Con A blasts were lysed at low to moderate levels, whereas no detectable lysis of B10.S (H-2^s) Con A blasts was observed (Fig. 5B). These results suggest that Ly-49W is an activating receptor that recognizes one or more H-2^k and H-2^d product(s).

Ly-49W recognizes H-2D^k and H-2D^d

The previous results suggest that there is an H-2^k and perhaps an H-2^d product or products recognized by Ly-49W. To determine which H-2^k class I product(s) is recognized by Ly-49W, we attempted to block Ly-49W-mediated lysis of B10.BR (H-2^k) Con A blasts with Abs that recognize either H-2K^k or H-2D^k. Soluble PA was used to bind the Ab Fc domain to prevent ADCC by the FcR-expressing RNK-16 cells. Neither the 11-4.1 Ab, which recognizes H-2K^k (35), nor the isotype control Ab B27 M1 blocks lysis of B10.BR Con A blasts (Fig. 6). In contrast, the 15-5-5S Ab, which recognizes H-2D^k (36), blocks lysis of B10.BR Con A blasts by both Ly-49W transfectant clones 7E8 and 10G5 in a dose-dependent manner (Fig. 6). These results indicate that Ly-49W recognizes H-2D^k.

Con A blast lysis results using congenic mice suggest that Ly-49W has a low to moderate level of recognition of an H-2^d molecule. The Ly-49W transfectant 7E8 lyses B10.D2 Con A blasts at a moderate level and was therefore used in additional experiments to determine which H-2^d product can be recognized by Ly-49W. We took two complementary approaches to address this issue. First, 7E8 lysis of Con A blasts generated from intra-MHC recombinant mouse strains B10.b/d (H-2K^b, H-2D^d) and B10.d/b (H-2K^d, H-2D^b) was compared with the B10 (H-2^b)-negative control. Con A-activated T cell blasts from the intra-MHC recombinant that expresses H-2D^d were lysed, whereas those from the recombinant expressing H-2K^d and the B10 strain were not (Fig. 7A). Second, we attempted to block lysis of B10.D2 Con A blasts by the 7E8 Ly-49W transfectant clone with two Abs that recognize H-2D^d: 34-5-8S, which recognizes the D^d α 1/ α 2 domains, and 34-2-12S, which recognizes the D^d α 3 domain. Both D^d-specific Abs blocked Ly-49W-mediated RNK lysis of the B10.D2 blasts (Fig. 7B), whereas the isotype control Ab or protein A had no effect. The 34-5-8S Ab blocking was more effective than 34-2-12S through the titration of these Abs, consistent with previous studies of Ly-49A interactions with H-2D^d (22). This suggests that Ly-49W interacts with H-2D^d in the α 1/ α 2 domains, as does Ly-49A. In addition, because the 34-2-12S Ab also blocks Ly-49W interaction with D^d, albeit less efficiently, the α 3 domain may also contribute to the interaction, or this Ab may sterically hinder Ly-49W interaction with the D^d α 1/ α 2 domains. In any case, these results indicate that Ly-49W recognizes H-2D^d.

Lysis of H-2^k-expressing targets by Ly-49W-transfected RNK cells is inhibited by Abs that recognize Ly-49W

The foregoing experiments demonstrated that Ly-49W-transfected RNK cells readily recognize H-2D^k-expressing targets, while untransfected RNK cells do not recognize them at all. To confirm that




FIGURE 4. Ly-49W expressed on RNK-16 cells mediates rADCC with the Cwy-3 Ab. RNK-16 cells (A) and the Ly-49W transfectant clones 2C4 (B), 10G5 (C), and 7E8 (D) were incubated with the indicated Abs at 20 μ g/ml or with medium for 15 min before addition of FcR-bearing YB2/0 target cells and a 4-h cytotoxicity assay. The data represent the mean of triplicate wells \pm SD.




FIGURE 5. Ly-49W activation of RNK-16-mediated lysis of Con A blasts is MHC dependent. RNK-16 and Ly-49W-transfected RNK-16 cells were tested as effectors in the lysis of Con A-activated T cell blast targets generated from splenocytes isolated from various mouse strains: A, H-2^k (AKR, CBA/J), H-2^d (B10.D2, DBA/2J), and H-2^b (C57BL/6); B, B10 congenic targets expressing H-2^k (B10.BR), H-2^d (B10.D2), and H-2^s (B10.S). E:T cell ratios are indicated and cytotoxicity was measured after 4 h. The data represent the mean of triplicate wells \pm SD.

the observed lytic activity is mediated by Ly-49W, we examined whether Abs recognizing Ly-49W could block the cytolytic activity. Both of the Ly-49W-recognizing Abs 4D11 and Cwy-3 reduced the killing of B10.BR Con A blasts by clone 10G5 (Fig. 8). The Cwy-3 Ab inhibited the majority of the lysis observed with the Ly-49W transfectant clone 10G5, and it was more effective than 4D11 (Fig. 8). Combining 4D11 and Cwy-3 Abs did not increase inhibition of lysis. Similar results were obtained with other Ly-49W transfectants (data not shown). These results support the role of Ly-49W in mediating RNK lysis of H-2D^k target cells.

Ly-49W^{NOD} transfectants of RNK-16 cells do not lyse NOD or NOR Con A blasts

The NOD and NOR mouse strains express H-2K^d and H-2D^b MHC molecules. Our previous results indicated that neither of

these MHC proteins is recognized by Ly-49W (Fig. 7A). To confirm that Ly-49W from NOD and NOR mice does not recognize self MHC, we compared lysis of Con A blasts generated from NOD, NOR, AKR, and B6 mice by Ly-49W RNK transfectants. As in previous experiments, AKR Con A blasts served as very good targets, while B6 blasts were not lysed by the Ly-49W-expressing RNK-16 effector cells (Fig. 9). In comparison, we found that NOD or NOR Con A blasts were not lysed by clones 2C4 and 10G5, and lysed to only a very limited extent, if at all, by the more highly cytolytic clone 7E8 (Fig. 9). These results demonstrate that Ly-49W from NOD and NOR mice does not recognize self Ags. We conclude from our studies that Ly-49W of NOD/NOR mice is an activating receptor specific for allogeneic class I MHC molecules.

Discussion

The B6 strain has served as a prototype for the study of Ly-49 gene expression and function. However, apparent strain-specific transcripts encoding novel activating receptors have recently been described: Ly-49P from the 129/J and NOD mouse strains (25, 40), and Ly-49L from CBA/J, C3H, and BALB/c mice (42) (Fig. 1B). Viable transcripts for none of these receptors have been found in the B6 mouse. We demonstrate in this study that NOD and NOR mice express two new mRNA transcripts encoding activating Ly-49 receptors. For one of these, Ly-49W, we could not find an allele in B6 mice. The other, Ly-49M, shares 98.7% nucleotide sequence identity with a transcript that we cloned from B6 mice and that incorporates the sequence of two genomic fragments previously designated Ly-49M. Interestingly, NOD Ly-49M transcripts appear to be functional, while B6 Ly-49M transcripts contain a premature stop codon. These findings present clear evidence that Ly-49 repertoires differ significantly between mouse strains. We further show that in at least one case, Ly-49M, this difference




FIGURE 6. Ly-49W recognizes H-2D^k. Lysis of B10.BR (H-2^k) Con A blasts by the 7E8 and 10G5 Ly-49W transfectants was determined in the presence of PA and various concentrations of 11-4.1 (anti-K^k), 15-5-5S (anti-D^k), or B27 M1 (isotype control) Abs. Abs were preincubated with PA (2 μ g/10 μ g of mAb) to prevent ADCC. Effector cells were added to provide a 12.5:1 E:T cell ratio. Cytotoxicity was measured after 4 h. Data are the means of triplicate wells \pm SD.




FIGURE 7. Lysis of H-2^d Con A blast targets by Ly-49W-transfected RNK cells is dependent on H-2^d recognition. *A*, Lysis by Ly-49W-transfected RNK-16 clone 7E8 of Con A blasts generated from intra-MHC recombinants B10.d/b (H-2K^d and D^b), B10.b/d (H-2K^b and D^d), and the B10 (H-2K^b and D^b) control strain. *B*, Lysis of B10.D2 Con A blasts by 7E8 in the presence of medium or PA in control wells (▨), or PA plus 34-5-8S (anti-D^d α 1/ α 2 domain, □), 34-2-12S (anti-D^d α 3 domain, ▩), or isotype control Ab B27 M1 (anti-HLA B27, ■) at the indicated concentrations. Abs were preincubated with PA (2 $\mu\text{g}/10 \mu\text{g}$ of Ab) to prevent ADCC. Abs and PA remained in the wells throughout the assay. The E:T ratio in *B* was 12.5:1. Cytotoxicity was measured after 4 h. Data are the means of triplicate wells \pm SD.

is due to a gene defect rather than to different levels of gene transcription.

Ly-49L, M, and W are closely related genes as indicated by sequence comparisons, suggesting that they result from relatively recent gene duplication and perhaps exon exchange. The ability of Ly-49 genes to undergo duplication and genetic recombination is similar to the functionally equivalent KIR receptors in primates, perhaps in response to similar evolutionary pressures (47–50). Evidence of human KIR gene recombination involving genes widely separated in the genome (47, 48) supports a contribution of gene conversion or nonhomologous recombination in the generation of receptor diversity. There are specific areas within the KIR gene complex in which these variations occur, suggesting that hot spots of gene duplication in the KIR gene complex may exist (48). The Ly-49 gene complex may have similar variability in different mouse strains, as the results in this study and others suggest (25, 40, 42). The relatively recent formation of Ly-49L, M, and W by gene duplication and exon exchange events suggest that these genes may be located near a similar hot spot in the NK gene complex region (51) encoding Ly-49 genes. Thus, gene duplication and




FIGURE 8. Lysis of H-2K^b-expressing targets by Ly-49W-transfected RNK-16 is inhibited by Abs that recognize Ly-49W. Cytotoxicity of B10.BR (H-2^k) Con A T cell blasts by the Ly-49W-transfected RNK-16 clone 10G5 was measured in the presence of medium or PA/PG alone (▨), or PA/PG with OX-8 (■), 4D11 (▩), Cwy-3 (□), or 4D11 plus Cwy-3 (▩), at the indicated concentrations. Abs were preincubated with PA/PG (4 $\mu\text{g}/10 \mu\text{g}$ Ab) for 30 min before addition to effector cells. Effector cells were incubated with the mAbs and PA/PG for 15 min before the cytotoxicity assay. The PA/PG and Abs remained in the wells throughout the assay. The E:T ratio was 12.5:1. Cytotoxicity was measured after 4 h. Data are the means of triplicate wells \pm SD.

nonhomologous recombination or gene conversion may contribute to the generation of polymorphism in both Ly-49 and KIR gene families.

In addition to substantial homology between Ly-49W, Ly-49M, and Ly-49L, there is a striking similarity of the Ly-49W CRD with that of the Ly-49G inhibitory receptor. For exons 5 to 7, encoding




FIGURE 9. Ly-49W^{NOD} RNK-16 transfectants do not lyse NOD or NOR Con A T cell blasts. RNK-16 and individual Ly-49W^{NOD} transfectant clones, 2C4, 7E8, and 10G5, were assayed for cytotoxicity against Con A T cell blasts from AKR/J, C57BL/6, NOD, and NOR mouse strains, at the indicated E:T cell ratios. Cytotoxicity was measured after 4 h. Data are the means of triplicate wells \pm SD.

the CRD, the amino acid sequence identity between Ly-49W and G is 97.6%. The sequence conservation drops to 83.8% for exon 4, which encodes the less well-conserved stalk region. For exons 2 and 3, the amino acid sequence identity is only 58.2% and, unlike Ly-49W and M, Ly-49G contains the ITIM motif in its cytoplasmic domain. Based on these results, Ly-49G and Ly-49W and M should be considered as inhibitory/activating pairs of Ly-49 receptors, similar to other examples such as: Ly-49A and Ly-49P (25, 40); Ly-49O and Ly-49D (40); Ly-49C/Ly-49I/Ly-49J and Ly-49H (41); NKG2-A and NKG2-C/NKG2-E (52, 53); NKR-P1B and NKR-P1A/NKR-P1C (54). The generation of inhibiting/activating receptor gene pairs may involve nonhomologous gene recombination events or gene conversion.

Ly-49W joins Ly-49D and Ly-49P as activating Ly-49 receptors that recognize class I MHC ligands (24, 25). Our findings with Ly-49W strengthen the conclusion that activating Ly-49 receptors, just like inhibitory Ly-49 receptors, recognize class I MHC ligands. We have demonstrated that Ly-49W is the first activating Ly-49 receptor to recognize an H-2^k product. Ligands for Ly-49W were determined to be H-2D^k and, with weaker interaction, H-2D^d. This novel recognition pattern contrasts with Ly-49D and Ly-49P, which only react significantly with H-2D^d (24, 25). From NK lytic assays using Con A blast target cells, it was determined that the inhibitory Ly-49G receptor of B6 mice recognizes D^d, D^r, and an undefined H-2^k product (55). This work is contradicted by the finding that Ly-49G does not bind soluble H-2D^k tetramers and does not mediate binding to H-2^k-bearing cells in cell-cell adhesion assays (56). However, the high sequence identity between the CRD of Ly-49G and Ly-49W and our finding that Ly-49W binds H-2D^k suggest that Ly-49G should be reexamined for H-2D^k reactivity.

With the availability of the Ly-49A/H-2D^d complex co-crystal structure (57), it is now possible to correlate sequence variation among Ly-49 and class I MHC molecules with their known interaction specificities. Ly-49A has a known binding specificity for H-2D^d with only weak binding to H-2D^k (22, 56). Conversely, we have demonstrated in this study that Ly-49W reacts strongly to D^k and only weakly to D^d. Among the D^d residues that interact directly with Ly-49A in the crystal structure, only two are not conserved in D^k (Lys¹⁷³-Asn¹⁷⁴ in D^d and Glu¹⁷³-Leu¹⁷⁴ in D^k). Similarly, among the Ly-49A residues that interact directly with D^d in the crystal structure, five positions are not conserved in Ly-49W, all in the hexapeptide 244–249 (NCDQVF in Ly-49A and DCGKSY in Ly-49W). Molecular modeling suggests that compensating changes within these sequences may confer the differential class I MHC specificities/affinities of Ly-49A and Ly-49W (not shown).

Self-tolerance of NK cells is believed to be maintained by expression and function of inhibitory receptors specific for self class I MHC proteins (10). Activating receptors may react with self MHC as well, and tolerance could be maintained by coexpression of inhibitory receptors that dominantly suppress activating receptor signals. For instance, it has been reported that tolerance of Ly-49D⁺ NK cells of self cells expressing H-2D^d is most likely maintained by the coexpression of Ly-49G and other Ly-49-inhibitory receptors (55). We demonstrate in this study that Ly-49W does not interact with self class I MHC proteins of the NOD/NOR mouse strains, but is instead alloreactive. For this reason, self-tolerance in NOD/NOR strains is not likely to be affected by Ly-49W. It remains to be determined whether, and if so how, Ly-49W⁺ NK cells are tolerant in F₁ animals such as CBA × NOD, in which an identified Ly-49W ligand is expressed.

We found that the Cwy-3 and 4D11 Abs were both reactive with Ly-49W, an activating receptor, and there were differences be-

tween the results obtained using Cwy-3 and 4D11. While rADCC using RNK-16 transfectants of Ly-49W could be demonstrated with the Cwy-3 Ab, the 4D11 Ab was unable to mediate rADCC (data not shown). Additionally, the Cwy-3 Ab was somewhat better at blocking Ly-49W-mediated lysis of H-2^k Con A blasts than 4D11 (Fig. 8). Differences in Ab affinity and/or site of attachment are most likely the cause of these different results. Our results suggest that it cannot be assumed that the inhibitory Ly-49G receptor is what is recognized by 4D11 in every mouse strain, because in addition, or instead, it may recognize an activating Ly-49 such as M, L, W, or related receptors in certain strains.

The existence of activating receptors that recognize class I MHC molecules is not readily predicted by the missing self hypothesis, which prompts the question: what is the role of MHC-specific activators in NK cell function? Activating Ly-49 members may function in conjunction with inhibitory receptors by recruiting kinases to phosphorylate ITIM sequences following ligand binding, thus augmenting the recruitment of SHP-1 and thereby the function of coexpressed inhibitory receptors. This possibility remains to be explored. The expression of inhibitory Ly-49 receptors occurs through a stochastic process that obeys a simple statistical product rule for expression of multiple Ly-49 receptors (10). In contrast, a recent report provides evidence to suggest that there is nonstochastic expression of two Ly-49 activators, D and H, in that there is a greater tendency for them to be coexpressed on NK cells (46). These observations suggest that Ly-49 activators may coordinate their activities and possibly function independent of inhibitory Ly-49 receptors in some NK cells (46). However, this analysis involved only two Ly-49-activating receptors, and it is now clear that several more exist. Further studies that include additional activating Ly-49 receptors will be necessary before a paradigm can be established for the function of this form of Ly-49 receptor. Identifying the extent and diversity of the Ly-49 family, both activating and inhibitory members, as well as their expression patterns and ligand specificities, should provide an opportunity to fully understand Ly-49 receptor functions. Identification and characterization of Ly-49W, a class I MHC-specific activating receptor, contribute to this goal.

Acknowledgments

We thank Dr. Mary Nakamura (University of California at San Francisco) for RNK-16 cells and for helpful advice in transfecting them, and Dr. Andrey Shaw (Washington University) for the BSRαEN expression vector. We also thank Dr. John Elliott (University of Alberta) for provision of NOD and NOR mice and COS-7 cells.

References

- Ortaldo, J. R., and D. L. Longo. 1988. Human natural lymphocyte effector cells: definition, analysis of activity, and clinical effectiveness. *J. Natl. Cancer Inst.* 80:999.
- Trinchieri, G. 1989. Biology of natural killer cells. *Adv. Immunol.* 47:187.
- Yu, Y. Y., V. Kumar, and M. Bennett. 1992. Murine natural killer cells and marrow graft rejection. *Annu. Rev. Immunol.* 10:189.
- George, T., Y. Y. Yu, J. Liu, C. Davenport, S. Lemieux, E. Stoneman, P. A. Mathew, V. Kumar, and M. Bennett. 1997. Allorecognition by murine natural killer cells: lysis of T-lymphoblasts and rejection of bone-marrow grafts. *Immunol. Rev.* 155:29.
- Ryan, J. C., E. C. Niemi, M. C. Nakamura, and W. E. Seaman. 1995. NKR-P1A is a target-specific receptor that activates natural killer cell cytotoxicity. *J. Exp. Med.* 181:1911.
- Moretta, A., C. Bottino, R. Millo, and R. Biassoni. 1999. HLA-specific and non-HLA-specific human NK receptors. *Curr. Top. Microbiol. Immunol.* 244:69.
- Bottino, C., R. Biassoni, R. Millo, L. Moretta, and A. Moretta. 2000. The human natural cytotoxicity receptors (NCR) that induce HLA class I-independent NK cell triggering. *Hum. Immunol.* 61:1.
- Long, E. O. 1999. Regulation of immune responses through inhibitory receptors. *Annu. Rev. Immunol.* 17:875.
- Ljunggren, H. G., and K. Karre. 1990. In search of the 'missing self': MHC molecules and NK cell recognition. *Immunol. Today* 11:237.
- Raulet, D. H. 1999. Development and tolerance of natural killer cells. *Curr. Opin. Immunol.* 11:129.

11. Lanier, L. L. 1998. NK cell receptors. *Annu. Rev. Immunol.* 16:359.
12. Vely, F., and E. Vivier. 1997. Conservation of structural features reveals the existence of a large family of inhibitory cell surface receptors and noninhibitory/activatory counterparts. *J. Immunol.* 159:2075.
13. Burshtyn, D. N., A. M. Scharenberg, N. Wagtmann, S. Rajagopalan, K. Berrada, T. Yi, J. P. Kinet, and E. O. Long. 1996. Recruitment of tyrosine phosphatase HCP by the killer cell inhibitor receptor. *Immunity* 4:77.
14. Nakamura, M. C., E. C. Niemi, M. J. Fisher, L. D. Shultz, W. E. Seaman, and J. C. Ryan. 1997. Mouse Ly-49A interrupts early signaling events in natural killer cell cytotoxicity and functionally associates with the SHP-1 tyrosine phosphatase. *J. Exp. Med.* 185:673.
15. Olcese, L., A. Cambiaggi, G. Semenzato, C. Bottino, A. Moretta, and E. Vivier. 1997. Human killer cell activatory receptors for MHC class I molecules are included in a multimeric complex expressed by natural killer cells. *J. Immunol.* 158:5083.
16. Tomasello, E., L. Olcese, F. Vely, C. Georgeon, M. Blery, A. Moqrigh, D. Gautheret, M. Djabali, M. G. Mattei, and E. Vivier. 1998. Gene structure, expression pattern, and biological activity of mouse killer cell activating receptor-associated protein (KARAP)/DAP-12. *J. Biol. Chem.* 273:34115.
17. Smith, K. M., J. Wu, A. B. Bakker, J. H. Phillips, and L. L. Lanier. 1998. Ly-49D and Ly-49H associate with mouse DAP12 and form activating receptors. *J. Immunol.* 161:7.
18. Brumbaugh, K. M., B. A. Binstadt, D. D. Billadeau, R. A. Schoon, C. J. Dick, R. M. Ten, and P. J. Leibson. 1997. Functional role for Syk tyrosine kinase in natural killer cell-mediated cytotoxicity. *J. Exp. Med.* 186:1965.
19. McVicar, D. W., L. S. Taylor, P. Gosselin, J. Willette-Brown, A. I. Mikhael, R. L. Geahlen, M. C. Nakamura, P. Linnemeyer, W. E. Seaman, S. K. Anderson, et al. 1998. DAP12-mediated signal transduction in natural killer cells: a dominant role for the Syk protein-tyrosine kinase. *J. Biol. Chem.* 273:32934.
20. Mason, L. H., J. Willette-Brown, A. T. Mason, D. McVicar, and J. R. Ortaldo. 2000. Interaction of Ly-49D⁺ NK cells with H-2D^d target cells leads to Dap-12 phosphorylation and IFN- γ secretion. *J. Immunol.* 164:603.
21. Karlhofer, F. M., R. K. Ribaldo, and W. M. Yokoyama. 1992. MHC class I alloantigen specificity of Ly-49⁺ IL-2-activated natural killer cells. *Nature* 358:66.
22. Kane, K. P. 1994. Ly-49 mediates EL4 lymphoma adhesion to isolated class I major histocompatibility complex molecules. *J. Exp. Med.* 179:1011.
23. Mason, L. H., J. R. Ortaldo, H. A. Young, V. Kumar, M. Bennett, and S. K. Anderson. 1995. Cloning and functional characteristics of murine large granular lymphocyte-1: a member of the Ly-49 gene family (Ly-49G2). *J. Exp. Med.* 182:293.
24. Nakamura, M. C., P. A. Linnemeyer, E. C. Niemi, L. H. Mason, J. R. Ortaldo, J. C. Ryan, and W. E. Seaman. 1999. Mouse Ly-49D recognizes H-2D^d and activates natural killer cell cytotoxicity. *J. Exp. Med.* 189:493.
25. Silver, E. T., D. E. Gong, C. S. Chang, A. Amrani, P. Santamaria, and K. P. Kane. 2000. Ly-49P activates NK-mediated lysis by recognizing H-2D^d. *J. Immunol.* 165:1771.
26. Delovitch, T. L., and B. Singh. 1997. The nonobese diabetic mouse as a model of autoimmune diabetes: immune dysregulation gets the NOD. *Immunity* 7:727.
27. Mason, L., S. L. Giardino, T. Hecht, J. Ortaldo, and B. J. Mathieson. 1988. LGL-1: a non-polymorphic antigen expressed on a major population of mouse natural killer cells. *J. Immunol.* 140:4403.
28. Chang, C. S., E. T. Silver, and K. P. Kane. 1999. Generation of a monoclonal antibody that recognizes a polymorphic epitope of C57BL/6 Ly-49G2. *Hybridoma* 18:423.
29. Springer, T. A. 1980. Cell surface differentiation in the mouse: characterization of "jumping" and "lineage" antigens using xenogeneic rat monoclonal antibodies. In *Monoclonal Antibodies*. R. H. Kennett, T. J. McKearn, and K. B. Bechtol, eds. Plenum Press, New York, p. 185.
30. Evans, G. A., D. H. Margulies, B. Shykind, J. G. Seidman, and K. Ozato. 1982. Exon shuffling: mapping polymorphic determinants on hybrid mouse transplantation antigens. *Nature* 300:755.
31. Ajitkumar, P., S. S. Geier, K. V. Kesari, F. Borriello, M. Nakagawa, J. A. Bluestone, M. A. Saper, D. C. Wiley, and S. G. Nathanson. 1988. Evidence that multiple residues on both the α -helices of the class I MHC molecule are simultaneously recognized by the T cell receptor. *Cell* 54:47.
32. Grumet, F. C., B. M. Fendly, and E. G. Engleman. 1981. Monoclonal anti-HLA-B27 antibody (B27M1): production and lack of detectable typing difference between patients with ankylosing spondylitis, Reiter's syndrome, and normal controls. *Lancet* 2:174.
33. Brodsky, F. M., P. Parham, C. J. Barnstable, M. J. Crumpton, and W. F. Bodmer. 1979. Monoclonal antibodies for analysis of the HLA system. *Immunol. Rev.* 47:3.
34. Thomas, M. L., and J. R. Green. 1983. Molecular nature of the W3/25 and MRC OX-8 marker antigens for rat T lymphocytes: comparisons with mouse and human antigens. *Eur. J. Immunol.* 13:855.
35. Oi, V. T., P. P. Jones, J. W. Goding, and L. A. Herzenberg. 1978. Properties of monoclonal antibodies to mouse Ig allotypes, H-2, and Ia antigens. *Curr. Top. Microbiol. Immunol.* 81:115.
36. Ozato, K., N. Mayer, and D. H. Sachs. 1980. Hybridoma cell lines secreting monoclonal antibodies to mouse H-2 and Ia antigens. *J. Immunol.* 124:533.
37. Reynolds, C. W., E. W. Bere, Jr., and J. M. Ward. 1984. Natural killer activity in the rat. III. Characterization of transplantable large granular lymphocyte (LGL) leukemias in the F344 rat. *J. Immunol.* 132:534.
38. Smith, H. R., F. M. Karlhofer, and W. M. Yokoyama. 1994. Ly-49 multigene family expressed by IL-2-activated NK cells. *J. Immunol.* 153:1068.
39. Ortaldo, J. R., A. T. Mason, R. Winkler-Pickett, A. Raziuddin, W. J. Murphy, and L. H. Mason. 1999. Ly-49 receptor expression and functional analysis in multiple mouse strains. *J. Leukocyte Biol.* 66:512.
40. Makrigiannis, A. P., P. Gosselin, L. H. Mason, L. S. Taylor, D. W. McVicar, J. R. Ortaldo, and S. K. Anderson. 1999. Cloning and characterization of a novel activating Ly49 closely related to Ly49A. *J. Immunol.* 163:4931.
41. Takei, F., J. Brennan, and D. L. Mager. 1997. The Ly-49 family: genes, proteins and recognition of class I MHC. *Immunol. Rev.* 155:67.
42. Makrigiannis, A. P., and S. K. Anderson. 2000. Ly-49 gene expression in different inbred mouse strains. *Immunol. Res.* 21:39.
43. McQueen, K. L., J. D. Freeman, F. Takei, and D. L. Mager. 1998. Localization of five new Ly49 genes, including three closely related to Ly49c. *Immunogenetics* 48:174.
44. Silver, E. T., J. F. Elliott, and K. P. Kane. 1996. Alternatively spliced Ly-49D and H transcripts are found in IL-2-activated NK cells. *Immunogenetics* 44:478.
45. Mason, L. H., S. K. Anderson, W. M. Yokoyama, H. R. Smith, R. Winkler-Pickett, and J. R. Ortaldo. 1996. The Ly-49D receptor activates murine natural killer cells. *J. Exp. Med.* 184:2119.
46. Smith, H. R., H. H. Chuang, L. L. Wang, M. Salcedo, J. W. Heusel, and W. M. Yokoyama. 2000. Nonstochastic coexpression of activation receptors on murine natural killer cells. *J. Exp. Med.* 191:1341.
47. Shilling, H. G., K. Lienert-Weidenbach, N. M. Valiante, M. Uhrberg, and P. Parham. 1998. Evidence for recombination as a mechanism for KIR diversification. *Immunogenetics* 48:413.
48. Wilson, M. J., M. Torkar, A. Haude, S. Milne, T. Jones, D. Sheer, S. Beck, and J. Trowsdale. 2000. Plasticity in the organization and sequences of human KIR/ILT gene families. *Proc. Natl. Acad. Sci. USA* 97:4778.
49. Martin, A. M., E. M. Freitas, C. S. Witt, and F. T. Christiansen. 2000. The genomic organization and evolution of the natural killer immunoglobulin-like receptor (KIR) gene cluster. *Immunogenetics* 51:268.
50. Khakoo, S. I., R. Rajalingam, B. P. Shum, K. Weidenbach, L. Flodin, D. G. Muir, F. Canavez, S. L. Cooper, N. M. Valiante, L. L. Lanier, and P. Parham. 2000. Rapid evolution of NK cell receptor systems demonstrated by comparison of chimpanzees and humans. *Immunity* 12:687.
51. Yokoyama, W. M., B. F. Daniels, W. E. Seaman, R. Hunziker, D. H. Margulies, and H. R. Smith. 1995. A family of murine NK cell receptors specific for target cell MHC class I molecules. *Semin. Immunol.* 7:89.
52. Lohwasser, S., P. Hande, D. L. Mager, and F. Takei. 1999. Cloning of murine NKG2A, B and C: second family of C-type lectin receptors on murine NK cells. *Eur. J. Immunol.* 29:755.
53. Silver, E. T., J. C. Lau, and K. P. Kane. 1999. Molecular cloning of mouse NKG2A and C. *Immunogenetics* 49:727.
54. Yokoyama, W. M., and W. E. Seaman. 1993. The Ly-49 and NKR-P1 gene families encoding lectin-like receptors on natural killer cells: the NK gene complex. *Annu. Rev. Immunol.* 11:613.
55. George, T. C., J. R. Ortaldo, S. Lemieux, V. Kumar, and M. Bennett. 1999. Tolerance and alloreactivity of the Ly49D subset of murine NK cells. *J. Immunol.* 163:1859.
56. Hanke, T., H. Takizawa, C. W. McMahon, D. H. Busch, E. G. Pamer, J. D. Miller, J. D. Altman, Y. Liu, D. Cado, F. A. Lemonnier, et al. 1999. Direct assessment of MHC class I binding by seven Ly49 inhibitory NK cell receptors. *Immunity* 11:67.
57. Tormo, J., K. Natarajan, D. H. Margulies, and R. A. Mariuzza. 1999. Crystal structure of a lectin-like natural killer cell receptor bound to its MHC class I ligand. *Nature* 402:623.